EPJ Web of Conferences will be set by the publisher
DOI: will be set by the publisher
© Owned by the authors, published by EDP Sciences, 2016

arXiv:1603.02067v1 [math.NA] 23 Feb 2016

Numerical Solution of a Nonlinear Integro-Differential Equation

Ján Buša[1,a], Michal Hnatič[2,3,4,b], Juha Honkonen[5,c], and Tomáš Lučivjanský[2,6]

[1] *Department of Mathematics and Theoretical Informatics, FEE&I, Technical University, Košice, Slovakia*
[2] *Faculty of Sciences, P. J. Šafarik University, Košice, Slovakia*
[3] *Institute of Experimental Physics SAS, Košice, Slovakia*
[4] *Bogoliubov Laboratory of Theoretical Physics, JINR, 141980 Dubna, Moscow Region, Russia*
[5] *Department of Military Technology, National Defence University, Helsinki, Finland*
[6] *Fakultät für Physik, Universität Duisburg-Essen, D-47048 Duisburg, Germany*

Abstract. An algorithm for the numerical solution of a nonlinear integro-differential equation arising in the single-species annihilation reaction $A + A \to \emptyset$ modeling is discussed. Finite difference method together with the linear approximation of the unknown function is considered. For divergent integrals presented in the equation for dimension $d = 2$ a regularization is used. Some numerical results are presented.

1 Introduction

The irreversible annihilation reaction $A + A \to \emptyset$ is a fundamental model of non-equilibrium physics. The reacting A particles are assumed to perform chaotic motion due to diffusion or some external advection field such as atmospheric eddy [1]. Many reactions of this type are observed in diverse chemical, biological or physical systems [2, 3].

In [1] the advection of reactive scalar using random velocity field generated by the stochastic Navier-Stokes equation, which is used for production a velocity field corresponding to thermal fluctuations [4, 5] and a turbulent velocity field with the Kolmogorov scaling behavior [6] is studied by three of the present authors, and the integro-differential equation for the number density is derived. No influence of the reactant on the velocity field itself is assumed.

In this paper we present some initial experiences with the numerical solution of the integro-differential equation mentioned above.

2 Problem Formulation

In [1] an integro-differential equation (72) for the mean number density $a(t)$ of chemically active molecules in anomalous kinetics of single-species annihilation reaction $A + A \to \emptyset$

$$\frac{\mathrm{d}a(t)}{\mathrm{d}t} = -2\lambda u\nu\mu^{-2\Delta} Z_4\, a^2(t) + 4\lambda^2 u^2\nu^2\mu^{-4\Delta} \int_0^t \frac{a^2(t')\,\mathrm{d}t'}{[8\pi u\nu(t-t')]^{d/2}}, \qquad a(0) = a_0 \tag{1}$$

[a] e-mail: jan.busa@tuke.sk
[b] e-mail: hnatic@saske.sk
[c] e-mail: juha.honkonen@helsinki.fi

is derived. The integral in Eq. (1) diverges at the upper limit t' in space dimensions $d \geq 2$.

We will study a numerical solution to the Initial Value Problem for the equation:

$$\frac{\mathrm{d}a(t)}{\mathrm{d}t} = -2\lambda D a^2(t) + 4\lambda^2 D^2 \int_0^t \frac{a^2(t')\,\mathrm{d}t'}{[8\pi D(t-t')]^{d/2}}, \qquad a(0) = a_0, \tag{2}$$

which corresponds to $D = u\nu$, $\Delta = 0$, and $Z_4 = 1$.

3 Case $d = 2$

For $d = 2$ the singularity in the integral on the right side of Eq. (2) at $t' = t$ is divergent. This is a consequence of the UV divergences in the model above the critical dimension $d_c = 2$, and near the critical dimension is remedied by the UV renormalization of the model [1]. In this paper we use another approach to overcome this problem. We will use the following *regularization*:

$$\frac{\mathrm{d}a}{\mathrm{d}t} = -2\lambda D a^2 + 4\lambda^2 D^2 \int_0^t \frac{a^2(t')\,\mathrm{d}t'}{8\pi\{D(t-t') + \ell^2\}}, \qquad a(0) = a_0. \tag{3}$$

We rewrite Eq. (3) to the form

$$\frac{\mathrm{d}a}{\mathrm{d}t} = -\alpha a^2 + \beta \int_0^t \frac{a^2(t')\,\mathrm{d}t'}{t - t' + \gamma} \tag{4}$$

with

$$\alpha = 2\lambda D, \qquad \beta = \frac{\alpha^2}{8\pi D}, \qquad \gamma = \frac{\ell^2}{D}.$$

Let us consider the Eq. (4). We will use the difference method to solve it numerically. We will consider time discretization with the time step Δt:

$$t_k = k \cdot \Delta t, \quad a(t_k) = a_k, \qquad k = 0, 1, 2, \ldots \tag{5}$$

For the discretization of the left side we will use two formula – the first and the second order finite difference approximations:

$$\left.\frac{\mathrm{d}a}{\mathrm{d}t}\right|_{t=t_k} \approx \frac{a_k - a_{k-1}}{\Delta t} \qquad \text{or} \qquad \left.\frac{\mathrm{d}a}{\mathrm{d}t}\right|_{t=t_k} \approx \frac{3a_k - 4a_{k-1} + a_{k-2}}{2\Delta t}. \tag{6}$$

For the right-side integral approximation we will use piecewise linear approximation of the function $a(u)$:

$$\int_0^{t_k} \frac{a^2(t')\,\mathrm{d}t'}{t_k - u + \gamma} = \sum_{i=0}^{k-1} \int_{i\cdot\Delta t}^{(i+1)\cdot\Delta t} \frac{a^2(t')\,\mathrm{d}t'}{t_k - t' + \gamma} \approx \tag{7}$$

$$\approx \sum_{i=0}^{k-1} \int_{i\cdot\Delta t}^{(i+1)\cdot\Delta t} \frac{[t'\cdot(a_{i+1} - a_i)/\Delta t + (i+1)\cdot a_i - i\cdot a_{i+1}]^2\ \mathrm{d}t'}{t_k - t' + \gamma}.$$

Integrals in the right side of Eq. (7) we calculate analytically using

$$\gamma = \delta \cdot \Delta t \tag{8}$$

in the following way

$$\int_{i\cdot\Delta t}^{(i+1)\cdot\Delta t} \frac{[t'\cdot(a_{i+1} - a_i)/\Delta t + (i+1)\cdot a_i - i\cdot a_{i+1}]^2\ \mathrm{d}t'}{t_k - t' + \gamma} =$$

$$= \int_{i\cdot\Delta t}^{(i+1)\cdot\Delta t} \frac{[(t' - k\cdot\Delta t - \delta\cdot\Delta t)\cdot(a_{i+1}-a_i)/\Delta t + (k+\delta)\cdot(a_{i+1}-a_i) + (i+1)\cdot a_i - i\cdot a_{i+1}]^2\ \mathrm{d}t'}{k\cdot\Delta t - t' + \delta\cdot\Delta t} =$$

$$= \int_{i\cdot\Delta t}^{(i+1)\cdot\Delta t} \frac{[(k+\delta-i)\cdot a_{i+1} - (k+\delta-i-1)a_i - (k\cdot\Delta t - t' + \delta\cdot\Delta t)\cdot(a_{i+1}-a_i)/\Delta t]^2\ \mathrm{d}t'}{k\cdot\Delta t - t' + \delta\cdot\Delta t} =$$

$$= [(k+\delta-i)a_{i+1} - (k+\delta-i-1)a_i]^2 \cdot \int_{i\cdot\Delta t}^{(i+1)\cdot\Delta t} \frac{\mathrm{d}t'}{k\cdot\Delta t - t' + \delta\cdot\Delta t} -$$

$$-\frac{2\,[(k+\delta-i)a_{i+1} - (k+\delta-i-1)a_i]\cdot(a_{i+1}-a_i)}{\Delta t} \int_{i\cdot\Delta t}^{(i+1)\cdot\Delta t} \frac{(k\cdot\Delta t - t' + \delta\cdot\Delta t)\,\mathrm{d}t'}{(k\cdot\Delta t - t' + \delta\cdot\Delta t)} +$$

$$+\frac{(a_{i+1}-a_i)^2}{(\Delta t)^2} \int_{i\cdot\Delta t}^{(i+1)\cdot\Delta t} \frac{(k\cdot\Delta t - t' + \delta\cdot\Delta t)^2\,\mathrm{d}t'}{(k\cdot\Delta t - t' + \delta\cdot\Delta t)} =$$

$$= [(k+\delta-i)a_{i+1} - (k+\delta-i-1)a_i]^2 \cdot \left[-\ln(k\cdot\Delta t - t' + \delta\cdot\Delta t)\right]_{i\cdot\Delta t}^{(i+1)\cdot\Delta t} -$$

$$-\frac{2\,[(k+\delta-i)a_{i+1} - (k+\delta-i-1)a_i]\,(a_{i+1}-a_i)}{\Delta t}\cdot\Delta t + \frac{(a_{i+1}-a_i)^2}{(\Delta t)^2}\cdot\left[\frac{(k\cdot\Delta t - t' + \delta\cdot\Delta t)^2}{-2}\right]_{i\cdot\Delta t}^{(i+1)\cdot\Delta t} =$$

$$= \left\{[(k+\delta-i)a_{i+1} - (k+\delta-i-1)a_i]^2 \cdot \ln\frac{k+\delta-i}{k+\delta-i-1} - \left(k+\delta-i+\frac{1}{2}\right)(a_{i+1}-a_i)^2 - 2a_i(a_{i+1}-a_i)\right\}, \quad (9)$$

$i = 0, 1, \ldots, k-1$.

Putting approximations Eq. (6), Eq. (7), and Eq. (9) into Eq. (4) we arrive at the quadratic equations Eq. (11) and Eq. (12) in more explicit form with respect to a_k:

$$0 = \frac{3a_k - 4a_{k-1} + a_{k-2}}{2\Delta t} + \alpha\cdot a_k^2 - \quad (10)$$

$$-\beta\cdot\sum_{i=0}^{k-2}\left\{[(k+\delta-i)a_{i+1} - (k+\delta-i-1)a_i]^2\cdot\ln\frac{k+\delta-i}{k+\delta-i-1} - \left(k+\delta-i+\frac{1}{2}\right)(a_{i+1}-a_i)^2 - 2a_i(a_{i+1}-a_i)\right\} -$$

$$-\beta\cdot\left\{[(1+\delta)a_k - \delta a_{k-1}]^2\cdot\ln\frac{1+\delta}{\delta} - \left(1+\delta+\frac{1}{2}\right)(a_k - a_{k-1})^2 - 2a_{k-1}(a_k - a_{k-1})\right\}$$

or in the standard form

$$\left[\alpha - \beta\cdot\left\{(1+\delta)^2\cdot\ln\frac{1+\delta}{\delta} - \left(\frac{3}{2}+\delta\right)\right\}\right]\cdot a_k^2 + \left[\frac{3}{2\Delta t} - \beta\cdot a_{k-1}\cdot\left\{2\delta + 1 - 2(1+\delta)\delta\cdot\ln\frac{1+\delta}{\delta}\right)\right\}\right]\cdot a_k -$$

$$-\frac{4a_{k-1} - a_{k-2}}{2\Delta t} - \beta\cdot a_{k-1}^2\cdot\left\{\frac{1}{2} - \delta + \delta^2\cdot\ln\frac{1+\delta}{\delta}\right\} - \quad (11)$$

$$-\beta\cdot\sum_{i=0}^{k-2}\left\{[(k+\delta-i)(a_{i+1}-a_i)+a_i]^2\cdot\ln\frac{k+\delta-i}{k+\delta-i-1} - (a_{i+1}-a_i)\left[(k+\delta-i)(a_{i+1}-a_i)+a_i+\frac{a_{i+1}+a_i}{2}\right]\right\} = 0.$$

For the first step we have

$$\frac{a_1 - a_0}{\Delta t} + \alpha a_1^2 - \beta\left\{[(1+\delta)a_1 - \delta a_0]^2\cdot\ln\frac{1+\delta}{\delta} - \left(1+\delta+\frac{1}{2}\right)(a_1 - a_0)^2 - 2a_0(a_1 - a_0)\right\} = 0$$



Figure 1. Logarithms of the number density $a(t)$ together with its fit (above), and the fitting error (below)

or in the standard form

$$\left[\alpha-\beta\cdot\left\{(1+\delta)^2\cdot\ln\frac{1+\delta}{\delta}-\left(\frac{3}{2}+\delta\right)\right\}\right]\cdot a_1^2+\left[\frac{1}{\Delta t}-\beta\cdot a_0\cdot\left\{2\delta+1-2(1+\delta)\delta\cdot\ln\frac{1+\delta}{\delta}\right)\right\}\right]\cdot a_1-$$

$$-\frac{a_0}{\Delta t}-\beta\cdot a_0^2\cdot\left\{\frac{1}{2}-\delta+\delta^2\cdot\ln\frac{1+\delta}{\delta}\right\}=0. \tag{12}$$

The algorithm consists of the successive calculation of the values a_k, $k = 1, 2, \ldots, K$ solving the quadratic equation Eq. (12) at the first step using the value $a_0 = a(0)$, and further solving Eqs. (11) using previously determined values a_i, $i = 0, \ldots, k-1$.

4 Numerical results

Below the results for $a(0) = 5000$, $\lambda = 0.1$, $D = 0.4$ and $\ell = 0.001$ are presented. All calculations are done with the uniform time step Δt.

Figure 1 shows the results on interval [0;10] for $\Delta t = 1/3200$. Above the logarithm of $a(t)$ is shown together with the logarithm of its approximation

$$a(t)\approx\frac{26.528}{(t-0.24389)^{0.5026}} \tag{13}$$

Table 1. Comparison of the a(t) values at selected points for different values Δt

Δt	$t = 0.01$	$t = 0.1$	$t = 1$	$t = 10$
0.01	2028.8975	130.40166	41.991715	12.607961
0.005	1338.5228	158.18202	35.781151	10.549856
0.0025	1077.5108	157.67282	32.302138	9.379055
0.00125	1062.3410	155.70709	30.581382	8.792275
0.000625	1067.8149	154.62549	29.841981	8.538087
0.0003125	1068.6433	154.18457	29.566128	8.442899



Figure 2. Precision order $p(t)$ for four cases: 0.01–0.005–0.0025, ..., 0.00125–0.000625–0.0003125

on interval [7.5;10]. Below the approximation error less than cca. 10^{-5} is shown.

Table 1 compares the results at selected points t for the "increasing precision" for the decreasing values Δt from 0.01 to 0.0003125. Even for the step $\Delta t = 1/3200 = 0.0003125$ the results may not be considered to have the sufficient precision. If the step number is increasing two times, the calculation time is increasing four times.

If we suppose, that the difference between the numerical value $a_{\Delta t}(t)$ for the step Δt and the "exact" value $a(t)$ has the order p, e.g.,

$$|a_{\Delta t}(t) - a(t)| \approx C \cdot (\Delta t)^p, \tag{14}$$

where both C and p depends on t, then using the successive approximations for different steps Δt, $\Delta t/2$, and $\Delta t/4$ we get

$$p(t) \approx \log_2 \left| \frac{a_{\Delta t}(t) - a_{\Delta t/2}(t)}{a_{\Delta t/2}(t) - a_{\Delta t/4}(t)} \right|. \tag{15}$$

Figure 2 shows the "precision order" $p(t)$ calculated for four grid sequences: 0.01–0.005–0.0025, 0.005–0.0025–0.00125, 0.0025–0.00125–0.000625, and 0.00125–0.000625–0.0003125. One can see, that for $t > 0.5$ the $p(t)$ behavior is smooth, and the orders are cca. 0.8, 1.0, 1.2, and 1.4, respectively. For smaller t-values the $p(t)$ behavior is more complicated.

It seems to be reasonable to study analytically the behavior of the function $a(t)$ for the small values of t, and start the numerical computation from some point $t > 0$. Also non-uniform grid could be considered.

5 Conclusions

Numerical results presented above could give us some basic imagination about the behavior of the number density function $a(t)$. However, further improvements of the algorithm are necessary.

It will be also interesting to try to solve the renormalized integro-differential equation

$$\frac{\mathrm{d}a(t)}{\mathrm{d}t} = -2\lambda u v \mu^{-2\Delta}\, a^2(t) + 2\lambda u v \mu^{-2\Delta} a^2(t) \left\{ \frac{\lambda}{4\pi} \left[\gamma + \ln(2u v \mu^2 t) \right] \right\} + \\ + \frac{\lambda^2 u v \mu^{-2\Delta}}{2\pi} \int_0^t \frac{[a^2(t') - a^2(t)]\,\mathrm{d}t'}{t - t'}, \tag{16}$$

where $\Delta = (d-2)/2$ and $\gamma \doteq 0.57721$ is Euler's constant, presented in [1], and compare the results.

Another possibility is to study a behavior of the solution of the problem (1) for $d \to 2^-$.

Acknowledgements

The work was supported by VEGA Grant 1/0222/13 of the Ministry of Education, Science, Research and Sport of the Slovak Republic.

References

[1] J. Honkonen, M. Hnatič and T. Lučivjanský, EPJ B, **86** : 214 (2013)
[2] B. Derrida, V. Hakim and V. Pasquier, Phys. Rev. Lett. **75**, 751 (1995)
[3] R. Kroon, H. Fleurent and R. Sprik, Phys. Rev. E **47**, 2462 (1993)
[4] D. Forster, D.R. Nelson and M.J. Stephen, Phys. Rev. Lett. **36**, 867 (1976)
[5] D. Forster, D.R. Nelson and M.J. Stephen, Phys. Rev. A **16**, 732 (1977)
[6] L.T. Adzhemyan, A.N. Vasil'ev and Y.M. Pis'mak, Teor. Mat. Fiz. **57**, 268 (1983)